

PARAPHYLY OF THE *ENOPLIGNATHA OVATA* GROUP (ARANEAE, THERIDIIDAE) BASED ON DNA SEQUENCES

A.-M. Tan¹, R.G. Gillespie¹ and G.S. Oxford²: ¹Center for Conservation Research and Training, University of Hawaii, 3050 Maile Way, Gilmore 409, Honolulu, Hawaii 96822, USA; ²Department of Biology, University of York, P.O. Box 373, York YO1 5YW, UK

ABSTRACT: Five species of *Enoplognatha* Pavesi 1880 were recently recognized as a monophyletic *Enoplognatha ovata* group based on morphological data. We analyzed the *E. ovata* clade for monophyly using four species in the *E. ovata* group (*E. ovata* (Clerck 1757), *E. latimana* Hippa & Oksala 1982, *E. margarita* Yaginuma 1964 and *E. afrodite* Hippa & Oksala 1983) and three other closely related taxa (*E. japonica* Bösenberg & Strand 1906, *E. thoracica* (Hahn 1833), and *E. intrepida* Sørensen 1898). Two species of the presumed sister genus (*Steatoda* Sundevall 1833) were employed as outgroups. The results indicate that the “*E. ovata* clade” is not monophyletic.

The genus *Enoplognatha* Pavesi 1880 is characterized by the presence of a large colulus, a plesiomorphic character for the family; and accordingly, the genus is generally considered one of the more primitive groups in the Theridiidae. The spiders are medium-to-large sized with a subspherical abdomen. Females have a tooth on the posterior margin of the chelicerae; males usually have enlarged chelicerae, with enlarged teeth on the posterior margin, and have the paracymbium on the margin of the cymbium. The genus is very close to *Steatoda* Sundevall 1833, medium-to-large sized spiders, again characterized by a very large colulus (Levi 1962; Levy & Amitai 1981). The chelicerae are often enlarged in males, and have one or more teeth on the anterior margin, none on the posterior margin.

Enoplognatha is well known because of the striking color and pattern polymorphism exhibited by representative species in the genus, which has been most intensively studied in *E. ovata* (Clerck 1757). Three distinct morphs have been described in *E. ovata* (Lockett & Millidge 1951; Hippa & Oksala 1979; Oxford 1976): *lineata* (all yellow), *redimita* (yellow with two dorsolateral carmine stripes on the abdomen), and *ovata* (yellow with a solid shield of carmine on the dorsal surface of the abdomen). The color pattern variation in *E. ovata* is genetically determined, and has been the subject of numerous studies on the genetics and evolution of the color polymorphism

(Hippa & Oksala 1979, 1981; Oxford 1983, 1985, 1989, 1991, 1992; Oxford & Reillo 1993; Reillo & Wise 1988a, b). Consistent with most invertebrate color polymorphisms (Haldane 1939) the dominance hierarchy of the expression of morphs in *E. ovata* follows the inverse of morph frequencies in nature, i.e., the least dominant (or most recessive) allele is most frequent; the most dominant is the rarest. For the mode of inheritance of the polymorphism in *E. ovata*, Oxford (1983) has proposed a two locus model: one locus is concerned with pattern and color, the other with the regulation of this color locus during development. When red-pigmented alleles are linked to the late developing allele, the color morphs are sex-limited: males are *lineata* no matter which allele they carry. *Enoplognatha latimana* Hippa & Oksala 1982 shares color, regulatory, and black spotting polymorphisms with *E. ovata* (Oxford 1992), although *E. latimana* lacks the *ovata* color morph.

In the 1980s Hippa & Oksala (1982) erected the *E. ovata* group to include *E. ovata sensu stricto*, *E. latimana*, and *E. penelope* Hippa & Oksala 1982. Members of the group share the following characters: trichobothrium on the first metatarsus subapical; elongated, sclerotized and subtubular tip of conductor in male palp; female vulva with massive copulatory pockets and abdomen with sharply delimited dorsolateral black spots (Hippa & Oksala 1982). Further examination of material

from Europe and Japan added another two species to the *E. ovata* group (Hipps & Oksala 1983): *E. afrodite* Hipps & Oksala 1983 and *E. margarita* Yaginuma 1964. *E. margarita* shares the subapical trichobothria and sclerotized subtubular tip of the conductor with *E. ovata*, *E. latimana* and *E. penelope*. However, it lacks the massive copulatory pockets. Considering all of these characters as synapomorphies, Hipps & Oksala (1983) hypothesized that *E. margarita* was the closest sister to (*E. ovata* + *E. latimana* + *E. penelope*). *Enoplognatha afrodite* has a similar body shape, ground color and spotting pattern to (*E. ovata* + *E. latimana* + *E. margarita*) but lacks these synapomorphies. Accordingly, Hipps & Oksala considered *E. afrodite* as the most ancestral species in the group.

More recently, Oxford & Reillo (1994) questioned the phylogeny of the *E. ovata* group proposed by Hipps & Oksala. Their concern arose because *E. ovata*, *E. latimana*, *E. penelope* and *E. afrodite* all have European distributions (although the former two have been introduced into North America). All occur in the Mediterranean region; but only *E. latimana* and *E. ovata* occur further north, with *E. ovata* alone extending well into northern Europe. Based on this distributional information, Oxford & Reillo hypothesized a possible Mediterranean origin of the *E. ovata* group, suggesting that the Asian *E. margarita* may have been phylogenetically misplaced by Hipps & Oksala. Indeed, the phylogeny presented by Hipps & Oksala was open to criticism because of the lack of a suitable outgroup for character polarization, few (only nine) characters used, and because there was no quantitative assessment of phylogeny.

In the current study we examined four species in the *E. ovata* group, and three other species of *Enoplognatha*: *E. japonica* Bösenberg & Strand 1906 from Japan, *E. thoracica* (Hahn 1833) from England, and *E. intrepida* Sørensen 1898 from North America. As outgroups in the analysis we used two species of *Steatoda*: *S. grossa* (C.L. Koch 1838) and *S. bipunctata* (Linnaeus 1758). We examined the pattern of sequence evolution in the *E. ovata* group to ascertain the monophyly of the clade. In this way we can evaluate the hypothesis that the Mediterranean served as the center of origin for the group as suggested by Oxford & Reillo (1994).

METHODS

Spiders sequenced.—*Enoplognatha*: *E. ovata*, two individuals from two localities: Grimes Graves, Norfolk, U.K., collected by G.S. Oxford, June 1991; and Berceto, Italy, collected by G.S. Oxford & P.R. Reillo, August 1991. *E. latimana*, one individual: Grimes Graves, Norfolk, U.K., collected by G.S. Oxford, June 1991. *E. afrodite*, one individual: near Carcassonne, S. France, collected by S. Peet, July 1988. *E. margarita*, one individual: Nukabira, Kamishihoro-cho, Hokkaido, Japan, collected by M. Matsuda, August 1992. Other *Enoplognatha* species examined: *E. japonica*, one individual: Hokkaido, Japan, collected by M. Matsuda, July 1989; *E. thoracica*, one individual: Flatford Mill, Suffolk, U.K., collected by C.J. Smith, May 1978; *E. intrepida*, one individual: Third Hill Mountain, Berkeley County, West Virginia, USA, collected by P.J. Martinat, May 1986 (det. D.T. Jennings, deposited in Smithsonian, Museum of Natural History). We also extracted DNA from *E. penelope*, one individual: Sami, Kefallinia, Greece, collected by J. Murphy, May 1987. However, we were not successful in amplifying the product. Outgroups: We used two species of *Steatoda* as the outgroup: *Steatoda grossa*: Molokai, Hawaii, collected by A.-M. Tan & G.S. Oxford October 1993 and *S. bipunctata*: Yorkshire, U.K., collected by G.S. Oxford, January 1994. Voucher specimens for all species used are at the Center for Conservation Research and Training, University of Hawaii.

DNA extraction and sequencing.—DNA samples were prepared by the conventional SDS-NaCl-Ethanol method (Medrano et al. 1990; Tan & Orrego 1992). Tissues from the legs or prosoma were placed in a 1.5 ml tube and ground with a pipette tip. After adding 15 μ l of proteinase K, the tissues were incubated at 55 °C overnight. Proteins were removed by salt precipitation. DNA was precipitated, washed in alcohol and preserved in 1 \times TE buffer (pH 8.0).

For both double and single stranded PCR amplification we used the following primers (Table 1): E and B2 for the less variable region of the 18S sequence; B and P for the more variable region of the 18S sequence; A and B2 for the 16S sequence. PCR amplification of double-stranded products was per-

Table 1.—Primers used. Position obtained refers to *Drosophila* (Clary & Wolstenholme 1985).

Gene primer	Primer sequence in <i>Drosophila</i>	Position obtained	# Base pairs	Reference
18S E	CTGGTTGATCTGCCAGTAG	24–553	529	modified from
18S B2	GCTGGCACCAGACTTGCCCTCC			Hillis & Dixon 1991
18S B	TTCCAGCTCCAATAGCGTAT	606–916	325	W.C. Wheeler & C. Hayashi,
18S P	GTCTTGCGACGGTCCAAGA			pers. comm.
16S A	CGCCTGTTTATCAAAAACAT	12864–13417	450	S.R. Palumbi & T. Hsiao,
16S B2	CTCCGGTTTGAACACTCAGATCA			pers. comm.

formed in 12.5 μ l volume with 38 cycles using *Thermus aquaticus* DNA polymerase (Saiki et al. 1985). Amplification was done with the following profile: 93 °C, 50 °C and 72 °C each for 30 seconds. Single strand products were prepared by asymmetric PCR (Gyllenstein & Erlich 1988) with 1:50 primer ratios in 50 μ l volumes and the same reaction profiles as above. The products were assessed by mini-gel electrophoresis using 5 μ l aliquots, and washed in sterilized distilled water with three cycles of dialysis using Millipore MC 30 (Amicon Corp.). Dideoxy chain termination sequencing (Sanger et al. 1977) was performed using the US Biochemicals Sequenase version 2.0 kit and ³⁵S labeled dATP. Negative controls were used in all PCR amplifications to make sure the sequences were not from contaminated sources. Sequences were confirmed by resequencing the same strand from another PCR product.

Phylogenetic analysis.—Ribosomal sequences were initially aligned using the program SeqEd 1.0.3 (Applied Biosystems 1995), after which alignment of multiple sequences was optimized in CLUSTAL W 1.4 (Higgins & Sharp 1988) in SeqPup 0.6 (Gilbert 1996). The entire first sequence is optimally aligned with the second entire sequence, with mismatches, gaps and insertions penalized equally, and with an additional gap length penalty for each residue in the insertion. Subsequent detailed alignment was by eye using the secondary structures (Kjer et al. 1994). The 18S sequences were aligned against the secondary structure of *Eurypelma californica* to match multiple sequences against conserved regions (Hendriks et al. 1988). The 16S sequences were aligned against *Drosophila yakuba* (Clary & Wolstenholme 1985), using the secondary structure of the region. Sequences were first analyzed using Maximum Likeli-

hood (ML) in PHYLIP (version 3.5c, Felsenstein 1993), using a generalized Jukes & Cantor (1969) model to allow for unequal base frequencies (Felsenstein 1981) as well as different rates of transitions and transversions. Sequences were also analyzed by Maximum Parsimony (MP) in PAUP (version 3.1.1, Swofford 1993). In both analyses gaps were treated as missing data. Bootstrap analyses (Felsenstein 1985) were used to estimate the statistical confidence of the different nodes in the trees.

RESULTS

The aligned sequences of the 18S region (Fig. 1) and 16S region (Fig. 2) are shown for each species (the two specimens of *E. ovata* were identical in sequence). Except for *E. thoracica* (18S only) and *E. intrepida* (16S only) we obtained 18S and 16S sequence for all species used. The data were first analyzed separately to determine the degree of congruence. The 18S sequences showed little bias in base composition, and no evidence for a transition: transversion (TS:TV) bias. The ML tree (using a TS:TV ratio of 1:1) was similar to the MP tree (using a branch-and-bound search) (Fig. 3A): (*E. thoracica* + *E. margarita*) and (*E. latimana* + *E. ovata*) were both discrete clades, and *E. japonica* fell outside all other species of *Enoplognatha*. The only difference between the analyses was that *E. afrodite* was placed with (*E. thoracica* + *E. margarita*) in the ML tree, while its position relative to (*E. thoracica* + *E. margarita*) and (*E. latimana* + *E. ovata*) was unresolved in the MP tree. Constraining *E. ovata*, *E. latimana*, *E. margarita* and *E. afrodite* to be monophyletic increased the length of the MP tree by two steps. We tested the monophyly of *E. ovata*, *E. latimana*, *E. margarita* and *E. afrodite* by calculating likelihood values (Fel-




Figure 4.—Phylogeny of representatives of the genus *Enoplognatha* based on Maximum Likelihood using the combined data set of 16S and 18S sequences. All branches are significant (approximate LRT, Felsenstein 1993). Parsimony analysis gave a similar topology but with less resolution: branches that were not supported by Maximum Parsimony are indicated as dashed lines; for branches that were supported, bootstrap values are given above nodes.

+ *E. ovata*) and (*E. japonica*, *E. intrepida* and *E. afrodite*) formed discrete clades. The primary difference between the analyses was that *E. margarita* was placed with (*E. japonica*, *E. intrepida* and *E. afrodite*) on the ML tree, but with (*E. latimana* + *E. ovata*) on the MP tree. Constraining *E. ovata*, *E. latimana*, *E. margarita* and *E. afrodite* to be monophyletic increased the length of the MP tree by six steps and resulted in a significantly lower log likelihood value for the ML tree (−1491.8 for the best tree, −1518.1 for the constrained tree).

Because the results from the two data sets were largely in agreement the data sets were combined and analyzed together. The resulting ML tree differed from the MP tree only in the degree of resolution it provided (Fig. 4). In all analyses *E. ovata* fell with *E. latimana*, *E. intrepida* with *E. japonica* (and in most cases with *E. afrodite*), *E. margarita* with *E. thoricica*. The *E. ovata* + *E. latimana* clade fell outside all others. We concluded that *E. ovata*, *E. latimana*, *E. margarita* and *E. afrodite* are not monophyletic, and again tested the robustness of these conclusions. Constraining *E. ovata*, *E. latimana*, *E. margarita* and *E. afrodite* to be monophyletic increased the length of the MP tree by three steps and gave a significantly lower log likelihood value for the ML tree (−3244.9 for the best tree, −3290.5 for the constrained tree).

DISCUSSION

The species *E. latimana*, *E. penelope*, *E. afrodite*, and *E. margarita* are similar in gross morphology to the well-studied *E. ovata*, and this similarity appears to be the basis for grouping these species into what has been considered to be a monophyletic clade (Hippha & Oksala 1983). The phylogenetic analysis presented here based on both the 16S and 18S sequences does not support monophyly of the “*E. ovata* group” as described by Hippha & Oksala (1983).

The *E. latimana* + *E. ovata* clade is strongly supported, and is consistent with evidence from color polymorphism: *E. ovata* and *E. latimana* share color, regulatory, and black spotting polymorphisms (Oxford 1992), although the latter species lacks the *ovata* color morph. These genetic traits suggest a recent common ancestor for this species pair. Color polymorphism has never been reported in any other species in the “*E. ovata* group”. However, the 18S and 16S data sets individually and combined consistently place *E. afrodite* and *E. margarita* outside the *E. latimana* + *E. ovata* clade, more closely associated with *E. japonica* and *E. intrepida*, and *E. thoricica* respectively. We have no molecular sequence data from *E. penelope*, and therefore cannot evaluate its position relative to others in the “*E. ovata* group”.

The results do not refute the Mediterranean center of origin hypothesis of Oxford & Reillo (1994), although the lack of monophyly of the group indicated by the current results demands a considerably larger representation from the genus be surveyed before their origin can be identified with any degree of certainty.

ACKNOWLEDGMENTS

The work reported here was supported in Hawaii by National Science Foundation Grant DEB 9207753 to R.G.G. and G.S.O., and in York by Natural Environment Research Council Grant GR9/1503 to G.S.O. For allowing destructive use of certain specimens, we would like to thank Jonathan Coddington and Scott Larcher (Smithsonian), Charles Griswold (California Academy), Herb Levi (MCZ, Harvard) and Norman Platnick (American Museum of Natural History). We thank M. Matsuda, J. Murphy, and the late C.J. Smith for providing us with specimens.

LITERATURE CITED

- Clary, D.O. & D.R. Wolstenholme. 1985. The mitochondrial DNA molecule of *Drosophila yakuba*: Nucleotide sequence, gene organization and genetic code. *J. Mol. Evol.*, 22:252–271.
- Felsenstein, J. 1981. Evolutionary trees from DNA sequences: a maximum likelihood approach. *J. Mol. Evol.*, 17:368–376.
- Felsenstein, J. 1985. Confidence limits on phylogenies: An approach using the bootstrap. *Evolution*, 39:783–791.
- Felsenstein, J. 1988. Phylogenies from molecular sequences: Inference and reliability. *Ann. Rev. Gen.*, 22:521–565.
- Felsenstein, J. 1993. PHYLIP. Phylogenetic Inference Package. Version 3.5c. University of Washington, Seattle.
- Gilbert, D.G. 1996. SeqPup, version 0.6., Biology Dept., Indiana University, Bloomington, Indiana 47405.
- Gyllenstein, V. & H. Erlich. 1988. Generation of single-stranded DNA by the polymerase chain reaction and its applications to direct sequencing of the HLA DQA locus. *Proc. Nat. Acad. Sci. USA*, 85:7652–7656.
- Haldane, J.B.S. 1939. The theory of the evolution of dominance. *J. Genet.*, 37:365–374.
- Hendriks, L., C. Van Broeckhoven, A. Vandenberghe, Y. Van De Peer & R. De Wachter. 1988. Primary and secondary structure of the 18S ribosomal RNA of the bird spider *Eurypelma californica* and evolutionary relationships among eukaryotic phyla. *European J. Biochem.*, 177: 15–20.
- Higgins, D.G., & P.M. Sharp. 1988. CLUSTAL: A package for performing multiple sequence alignment on a microcomputer. *Gene*, 73:237–244.
- Hillis, D.M. & M.T. Dixon. 1991. Ribosomal DNA: Molecular evolution and phylogenetic inference. *Quart. Rev. Biol.*, 66:411–453.
- Hippa, H. & I. Oksala. 1979. Colour polymorphism of *Enoplognatha ovata* (Clerck) (Araneae, Theridiidae) in western Europe. *Hereditas*, 90: 203–212.
- Hippa, H. & I. Oksala. 1981. Polymorphism and reproductive strategies of *Enoplognatha ovata* (Clerck) (Araneae, Theridiidae) in northern Europe. *Ann. Zool. Fennici*, 18:179–190.
- Hippa, H. & I. Oksala. 1982. Definition and revision of the *Enoplognatha ovata* (Clerck) group (Araneae: Theridiidae). *Entomol. Scandinavica*, 13:213–222.
- Hippa, H. & I. Oksala. 1983. Cladogenesis of the *Enoplognatha ovata* group (Araneae, Theridiidae), with description of a new Mediterranean species. *Ann. Ent. Fennici*, 49:71–74.
- Jukes, T.H. & C.R. Cantor. 1969. Evolution of protein molecules. Pp. 21–132, *In* Mammalian Protein Metabolism. (H.N. Munro, ed.). Academic Press, New York.
- Kishino, H. & M. Hasegawa. 1989. Evaluation of the maximum likelihood estimate of the evolutionary tree topologies from DNA sequence data, and the branching order in Hominoidea. *J. Mol. Evol.*, 29:170–179.
- Kjer, K.M., G.D. Baldrige & A.M. Fallon. 1994. Mosquito large subunit ribosomal RNA: Simultaneous alignment of primary and secondary structure. *Biochim. Biophys. Acta*, 1217:147–155.
- Levi, H.W. 1962. The spider genera *Steatoda* and *Enoplognatha* in America (Araneae, Theridiidae). *Psyche*, 69:11–36.
- Levy, G. & P. Amitai. 1981. The spider genus *Enoplognatha* (Araneae: Theridiidae) in Israel. *Zool. J. Linn. Soc.*, 72:43–67.
- Locket, G.H. & A.F. Millidge. 1951. *British Spiders*, Vol. 1. Ray Soc., London.
- Medrano, J.F., E. Aasen & L. Sharrow. 1990. DNA extraction from nucleated red blood cells. *Biotechniques*, 8:43.
- Oxford, G.S. 1976. The colour polymorphism in *Enoplognatha ovatum* (Clerck) (Araneae: Theridiidae)—temporal stability and spatial variability. *Heredity*, 36:369–381.
- Oxford, G.S. 1983. Genetics of colour and its regulation during development in the spider *Enoplognatha ovatum* (Clerck) (Araneae: Theridiidae). *Heredity*, 51:621–634.
- Oxford, G.S. 1985. Geographical distribution of phenotypes regulating pigmentation in the spider *Enoplognatha ovata* (Clerck) (Araneae: Theridiidae). *Heredity*, 55:37–45.
- Oxford, G.S. 1989. Genetics and distribution of black spotting in *Enoplognatha ovata* (Araneae: Theridiidae), and the role of intermittent drift in population differentiation. *Biol. J. Linn. Soc.*, 36: 111–128.
- Oxford, G.S. 1991. Visible morph-frequency variation in allopatric and sympatric populations of two species of *Enoplognatha* (Araneae: Theridiidae). *Heredity*, 67:317–324.
- Oxford, G.S. 1992. *Enoplognatha ovata* and *E. latimana*: a comparison of their phenologies and genetics in Norfolk populations. *Bull. British Arachnol. Soc.*, 9:13–18.
- Oxford, G.S. & P.R. Reillo. 1993. Trans-continental visible morph-frequency variation at homologous loci in two species of spider, *Enoplognatha ovata* s.s. & *E. latimana*. *Biol. J. Linn. Soc.*, 50:235–253.
- Oxford, G.S. & P.R. Reillo. 1994. The world distributions of species within the *Enoplognatha ovata* group (Araneae: Theridiidae): Implications for their evolution and for previous research. *Bull. British Arachnol. Soc.*, 9:226–232.
- Reillo, P.R. & D.H. Wise. 1988a. An experimental

- evaluation of selection on color morphs of the polymorphic spider *Enoplognatha ovata* (Araneae: Theridiidae). *Evolution*, 42:1172-1189.
- Reillo, P.R. & D.H. Wise. 1988b. Genetics of color expression in the spider *Enoplognatha ovata* (Araneae: Theridiidae) from coastal Maine. *American Midl. Nat.*, 119:318-326.
- Saiki, R.K., S. Scharf, F. Faloona, K.B. Mullis, G.T. Horn, H.A. Erlich & N. Arnheim. 1985. Enzymatic amplification of B-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science*, 230:1350-1354.
- Sanger, F., S. Nicklen & A.R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA*, 74:5463-5467.
- Swofford, D.L. 1993. PAUP: Phylogenetic analysis using parsimony, version 3.1.1. Smithsonian Institution, Washington, DC.
- Tan, A.-M. & C. Orrego. 1992. DNA amplification from museum collections of extracts originally intended for allozyme analysis. *Mol. Ecol.*, 1:95-97.

Manuscript received 10 February 1996, revised 1 October 1998.